

# Tourism Heritage and Cultural Studies

Volume: 4 Issue: 1  
(January-March), 2024  
(ISSN: 2791-2620)



## A Comprehensive Study on Post-Covid Complications among Patients in Vizianagaram District, Andhra Pradesh, India

Koneru Neha<sup>1</sup>, Mani Gudivada<sup>1</sup>

<sup>1</sup>Department of Zoology, Andhra University, Visakhapatnam, Andhra Pradesh, India

### ABSTRACT

This study was conducted on the analysis of the post-COVID complications of the patients in the three hospitals of Vizianagaram district in Andhra Pradesh, India. Over a period of four months from August 2022 to December 2022, data was collected to evaluate the occurrence of post-COVID complications by the intake of Mercury [in seafood], Zinc, and Iron supplements in the medicine. The study is first- ever documented in this specific region. A wide range of data was collected from three major hospitals in the Vizianagaram district to analyse the occurrence of post-COVID complications. This includes factors such as the Intake of zinc and intake of iron through medicine and, the intake of seafood in diet among the diabetic patients to increase the immune system. It was found that intake of seafood, zinc, and iron supplements, may lead to several post-COVID complications in diabetic patients. Based on the findings, it is recommended that most diabetic patients should take special care in their diet and intake of medicine to prevent such health complications.

Keywords: COVID-19, COVID associated Mucormycosis, Pneumococcal pneumonia, CAP, COPD, COVID Pneumonia, Dementia, Allergy, Gastro intestinal problems, Mercury, Iron, Zinc, Corticoides

**Correspondence to Koneru Neha**, Email: [nehakoneru.rs@andhrauniversity.edu.in](mailto:nehakoneru.rs@andhrauniversity.edu.in)

**Received:** January 11, 2024; **Accepted:** February 23, 2024; **Published:** March 30, 2024

**Citation:** Neha, K., & Gudivada, M. (2024). A comprehensive study on post covid complications among patients in vizianagaram district, Andhra Pradesh, India. *Tourism Heritage and Cultural Studies*, 4(1), 687-695. <https://doi.org/10.5281/zenodo.11066017>

**Copyright:** © 2024 by the authors. Licensee KMF Publishers ([www.kmf-publishers.com](http://www.kmf-publishers.com)). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## INTRODUCTION

After more than a year of the COVID-19 pandemic, many aspects of this new and complicated disease are still poorly understood and characterized [5,6], such as the frequency and risk factors associated with

complications after acute COVID-19 illness. Post-COVID-19 syndrome is defined by the symptoms that appear while or after suffering COVID-19 for more than 12 weeks [1,2,3,4]. The syndrome includes affection of respiratory, cardiovascular, neurological, gastrointestinal, and musculoskeletal systems,

psychological issues, ear, nose, throat, and dermatological symptoms [2,3,4,7,10,11,12]. The frequency of complications after the acute phase has important variations in the ranges from 10% to 93% [1,3,8,19,20,21,22,23,24,25]. The studies of post-COVID-19 complications are heterogeneous; including hospitalized versus non-hospitalized patients, patients' previous state of health, and severity of the disease. In this situation, accruing evidence on the frequency and the manifestation of post-COVID complications and related risk factors is needed


The present study aimed to describe post-COVID-19 complications, recovery, and return to the usual state of health in patients who suffered a SARS-CoV-2 infection and estimate the prevalence of the association of diverse risk factors with COVID-19.

## METHODOLOGY


In March 2022 a COVID-19 outbreak took place. An epidemiology questionnaire survey has been done to observe the outbreak of post-COVID complications in patients already suffered 3-6 months prior from COVID-19. There are some patients suffering from Diabetes. The observed post-COVID complications are pneumonia, Mucormycosis, cough, fatigue, joint pain, muscular pain, fever, sleep disturbances, dementia, headache, diarrhea, hair loss, and skin rashes.

## EPIDEMIOLOGY


The findings from studies reporting outcomes in subacute/ongoing symptomatic COVID-19 and chronic/post-COVID-19 complications are summarized in Table: 1. An observational cohort study from 3 hospitals in Vizianagaram district, Andhra Pradesh, India evaluated the outcomes of 120 patients discharged alive at 60d by utilizing medical record abstraction and telephone surveys (hereby referred to as the post-COVID-19 study). During the study period, 16% of patients died, while 14% of patients required re-admission. Of 100 patients who completed the telephone survey in this study, 83.3% of patients reported persistent symptoms, including 18.9% with new or worsened symptoms. Dyspnea while walking up the stairs (35.3%) was most commonly reported, while other symptoms included cough (7%), pneumonia (4%), mucormycosis (9.3%), joint pain (35.3%), chest pain (2.3%), sleep disturbance (49.3%), headache (24.65), diarrhea (18.3%), hair loss (71.3%) were the most commonly reported symptoms, with 55% of patients continuing to experience three or more symptoms. Fatigue, dyspnea, and psychological distress, such as post-traumatic stress disorder (PTSD), anxiety, depression, and concentration and sleep abnormalities, were noted in approximately 30% or more study participants at the time of follow-up.




Pie Chart-1: Data showing the patients suffered from POST- COVID complications in three different Hospitals, Vizianagaram district, Andhra Pradesh, India



Pie Chart-2: Data showing the patients suffered from POST- COVID complications in Hospital-1, Vizianagaram district, Andhra Pradesh, India



Pie Chart-3: Data showing the patients suffered from POST- COVID complications in Hospital-2, Vizianagaram district, Andhra Pradesh, India



Pie Chart-4: Data showing the patients suffered from POST- COVID complications in Hospital-3, Vizianagaram district, Andhra Pradesh, India

Table: 1- Findings from clinical studies on the prevalence of post-COVID complications in patients among three hospitals from Vizianagaram district, Andhra Pradesh, India

Site	Hospital-1	Hospital-2	Hospital-3
<b>Number of participants</b>	31	19	50
<b>FOLLOW UP</b>			
<b>Duration</b>	2-4 months post COVID symptoms onset	2-4 months post COVID symptoms onset	2-4 months post COVID symptoms onset
<b>Mode of follow up evaluation</b>	Telephone survey	Telephone survey	Telephone survey
<b>BASE LINE CHARACTERSTICS</b>			
<b>Age (yrs)</b>	20-55	20-55	20-55
<b>Female</b>	45%	52%	40%
<b>PRE HEALTH HISTORY</b>			
<b>Diabetes</b>	32%	42%	64%
<b>GENERAL SEQUELAE</b>			
<b>Fatigue</b>	-	-	-
<b>Joint pain</b>	25%	31%	50%
<b>Muscular pain</b>	25%	31%	50%
<b>Fever</b>	9%	5%	14%
<b>RESPIRATORY SEQUELAE</b>			
<b>Mucomycosis</b>	12%	0	16%
<b>Cough</b>	9%	0	6%
<b>Pneumonia</b>	6%	0	6%
<b>CARDIO VASCULAR SEQUELAE</b>			
<b>Chest pain</b>	3%	0	4%
<b>NEUROLOGICAL SEQUELAE</b>			
<b>Sleep disturbances</b>	38%	36%	74%
<b>Dementia</b>	NA	NA	NA
<b>Headache</b>	32%	42%	44%
<b>GASTROINTESTINAL SEQUELENCE</b>			
<b>Diarrhae</b>	6%	5%	6%
<b>DERMATOLOGICAL SEQUELENCE</b>			
<b>Hair loss</b>	83%	57%	74%
<b>Skin rashes</b>	-	-	-

## NUTRITIONAL CONSIDERATIONS

Severe COVID-19, similar to other critical illnesses, causes catabolic muscle wasting, feeding difficulties, and frailty, each of which is associated with an increased likelihood of poor outcomes [41]. Malnutrition has been noted in 26–45% of patients with COVID-19, as evaluated by the Malnutrition Universal Screening Tool in an Italian study [40]. Protocols to provide nutritional support for patients (many of whom suffered from respiratory distress, nausea, and diarrhea, with a resultant reduction in food intake) continue to be refined [41]. All post-acute COVID-19 follow-up studies that incorporated assessments of health-related quality of life and functional capacity measures have universally reported significant deficits in these domains, including at 6months in the post-acute COVID-19 Chinese study [42,43,44]. Model COVID-19 rehabilitation units such as those in Italy are already routinely assessing acute COVID-19 survivors for swallowing function, nutritional status, and measures of functional independence [40].

## CONCLUSION

The present survey concludes that, several post-COVID complications have arisen due to impairment of the immune system and also poor nutritional practices among the individuals suffering from COVID-19, in which females in the age group of 20-55 years are more prone to these infections.

Moreover, it is clear that care for patients with COVID-19 does not conclude at the time of hospital discharge, and interdisciplinary cooperation is needed for comprehensive care of these patients in the outpatient setting. As such, it is crucial for healthcare systems and hospitals to recognize the need to establish dedicated COVID19 clinics [40], where specialists from multiple disciplines are able to provide integrated care. Prioritization of follow-up care may be considered for those at high risk for post- COVID-19, including those

who had severe illness during COVID-19 and/or required care in an ICU, those most susceptible to complications and those with the highest burden of persistent symptoms. Given the global scale of this pandemic, it is apparent that the healthcare needs for patients with sequelae of COVID-19 will continue to increase for the foreseeable future. Rising to this challenge will require to control and make use of existing outpatient infrastructure, the development of scalable healthcare models and integration across disciplines for improved mental and physical health of survivors of COVID-19 in the long term.

## REFERENCES

1. Shah, W.; Hillman, T.; Playford, E.D.; Hishmeh, L. Managing the long term effects of COVID-19: Summary of NICE, SIGN, and RCGP rapid guideline. *BMJ* 2021, 372, n136. [Google Scholar] [CrossRef]
2. National Institute for Health and Care Excellence. COVID-19 Rapid Guideline: Managing the Long-Term Effects of COVID-19. 2020. Available online: <https://www.nice.org.uk/guidance/ng188>.
3. Greenhalgh, T.; Knight, M.; A’Court, C.; Buxton, M.; Husain, L. Management of post-acute COVID-19 in primary care. *BMJ* 2020, 370, m3026. [Google Scholar] [CrossRef]
4. Shanbehzadeh, S.; Tavahomi, M.; Zanjari, N.; Ebrahimi-Takanjani, I.; Amiri-Arimi, S. Physical and mental health complications post-COVID-19: Scoping review. *J. Psychosom. Res.* 2021, 147, 110525. [Google Scholar] [CrossRef]
5. Sridhar, S.; Nicholls, J. Pathophysiology of infection with SARS-CoV-2-What is known and what remains a mystery. *Respirology* 2021, 26, 652–665. [Google Scholar] [CrossRef] [PubMed]
6. Van Damme, W.; Dahake, R.; Delamou, A.; Ingelbeen, B.; Wouters, E.; Vanham, G.; van de Pas, R.; Dossou, J.P.; Ir, P.; Abimbola, S.; et al. The COVID-19 pandemic: Diverse

- contexts; different epidemics-how and why? *BMJ Glob. Health* 2020, 5, e003098. [Google Scholar] [CrossRef]
7. Salamanna, F.; Veronesi, F.; Martini, L.; Landini, M.P.; Fini, M. Post-COVID-19 Syndrome: The persistent symptoms at the post-viral stage of the disease: A systematic review of the current data. *Front. Med. Lausanne* 2021, 8, 653516. [Google Scholar] [CrossRef] [PubMed]
  8. Cairoli, E. ¿De qué hablamos cuando hablamos de pos-COVID-19? *Rev. Clin. Esp.* 2021, 221, 614–616. [Google Scholar] [CrossRef] [PubMed]
  9. Carod-Artal, F.J. Post-COVID-19 syndrome: Epidemiology, diagnostic criteria and pathogenic mechanisms involved. *Rev. Neurol.* 2021, 72, 384–396. [Google Scholar]
  10. Shah, W.; Hillman, T.; Playford, E.D.; Hishmeh, L. Managing the long term effects of COVID-19: Summary of NICE, SIGN, and RCGP rapid guideline. *BMJ* 2021, 372, n136. [Google Scholar] [CrossRef]
  11. National Institute for Health and Care Excellence. COVID-19 Rapid Guideline: Managing the Long-Term Effects of COVID-19. 2020. Available online: <https://www.nice.org.uk/guidance/ng188>
  12. Greenhalgh, T.; Knight, M.; A’Court, C.; Buxton, M.; Husain, L. Management of post-acute COVID-19 in primary care. *BMJ* 2020, 370, m3026. [Google Scholar] [CrossRef]
  13. Shanbehzadeh, S.; Tavahomi, M.; Zanjari, N.; Ebrahimi-Takanjani, I.; Amiri-Arimi, S. Physical and mental health complications post-COVID-19: Scoping review. *J. Psychosom. Res.* 2021, 147, 110525. [Google Scholar] [CrossRef]
  14. Sonnweber, T.; Sahanic, S.; Pizzini, A.; Luger, A.; Schwabl, C.; Sonnweber, B.; Kurz, K.; Koppelstätter, S.; Haschka, D.; Petzer, V.; et al. Cardiopulmonary recovery after COVID-19: An observational prospective multicentre trial. *Eur. Respir. J.* 2021, 57, 2003481. [Google Scholar] [CrossRef]
  15. Starace, M.; Iorizzo, M.; Sechi, A.; Alessandrini, A.M.; Carpanese, M.; Bruni, F.; Vara, G.; Apalla, Z.; Asz-Sigall, D.; Barruscotti, S.; et al. Trichodynia and telogen effluvium in COVID-19 patients: Results of an international expert opinion survey on diagnosis and management. *JAAD Int.* 2021, 5, 11–18. [Google Scholar] [CrossRef]
  16. Zarei, M.; Bose, D.; Nouri-Vaskeh, M.; Tajiknia, V.; Zand, R.; Ghasemi, M. Long-term side effects and lingering symptoms post COVID-19 recovery. *Rev. Med. Virol.* 2021, e2289. [Google Scholar] [CrossRef]
  17. Cozzi, G.; Amaddeo, A.; Barbi, E. Post-COVID syndrome: Turning convalescence into illness? *Lancet Reg. Health Eur.* 2021, 7, 100163. [Google Scholar] [CrossRef] [PubMed]
  18. Soriano, J.B.; Murthy, S.; Marshall, J.C.; Relan, P.; Diaz, J.V.; WHO Clinical Case Definition Working Group on Post-COVID-19 Condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect. Dis.* 2021. [Google Scholar] [CrossRef]
  19. Akbarialiabad, H.; Taghrir, M.H.; Abdollahi, A.; Ghahramani, N.; Kumar, M.; Paydar, S.; Razani, B.; Mwangi, J.; Asadi-Pooya, A.A.; Malekmakan, L.; et al. Long COVID, a comprehensive systematic scoping review. *Infection* 2021, 49, 1163–1186. [Google Scholar] [CrossRef]
  20. Amenta, E.M.; Spallone, A.; Rodriguez-Barradas, M.C.; El Sahly, H.M.; Atmar, R.L.; Kulkarni, P.A. Postacute COVID-19: An overview and approach to classification. *Open Forum Infect. Dis.* 2020, 7, ofaa509. [Google Scholar] [CrossRef] [PubMed]
  21. Miller, A. COVID-19: Not just an acute illness. *Trends Urol. Men’s Health* 2020, 11, 17–19. [Google Scholar] [CrossRef]



22. Poenaru, S.; Abdallah, S.J.; Corrales-Medina, V.; Cowan, J. COVID-19 and post-infectious myalgic encephalomyelitis/chronic fatigue syndrome: A narrative review. *Ther. Adv. Infect. Dis.* 2021, 8, 20499361211009385. [Google Scholar] [CrossRef] [PubMed]
23. Sigfrid, L.; Drake, T.M.; Pauley, E.; Jesudason, E.C.; Olliaro, P.; Lim, W.S.; Gillesen, A.; Berry, C.; Lowe, D.J.; McPeake, J.; et al. Long covid in adults discharged from UK hospitals after COVID-19: A prospective, multicentre cohort study using the ISARIC WHO Clinical Characterisation Protocol. *Lancet Reg. Health Eur.* 2021, 8, 100186. [Google Scholar] [CrossRef]
24. Nasserie, T.; Hittle, M.; Goodman, S.N. Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: A systematic review. *JAMA Netw. Open* 2021, 4, e2111417. [Google Scholar] [CrossRef]
25. Iqbal, F.M.; Lam, K.; Sounderajah, V.; Clarke, J.M.; Ashrafian, H.; Darzi, A. Characteristics and predictors of acute and chronic post-COVID syndrome: A systematic review and meta-analysis. *EClinicalMedicine* 2021, 36, 100899. [Google Scholar] [CrossRef]
26. Yong, S.J. Long COVID or post-COVID-19 syndrome: Putative pathophysiology, risk factors, and treatments. *Infect. Dis.* 2021, 53, 737-754. [Google Scholar] [CrossRef] [PubMed]
27. Carfi, A.; Bernabei, R.; Landi, F.; Gemelli against COVID-19 Post-Acute Care Study Group. Persistent Symptoms in Patients after Acute COVID-19. *JAMA* 2020, 324, 603-605. [Google Scholar] [CrossRef] [PubMed]
28. Osikomaiya, B.; Erinoso, O.; Wright, K.O.; Odusola, A.O.; Thomas, B.; Adeyemi, O.; Bowale, A.; Adejumo, O.; Falana, A.; Abdus-Salam, I.; et al. 'Long COVID': Persistent COVID-19 symptoms in survivors managed in Lagos State, Nigeria. *BMC Infect. Dis.* 2021, 21, 304. [Google Scholar] [CrossRef] [PubMed]
29. Peghin, M.; Palese, A.; Venturini, M.; De Martino, M.; Gerussi, V.; Graziano, E.; Bontempo, G.; Marrella, F.; Tommasini, A.; Fabris, M.; et al. Post-COVID-19 symptoms 6 months after acute infection among hospitalized and non-hospitalized patients. *Clin. Microbiol. Infect.* 2021, 27, 1507-1513. [Google Scholar] [CrossRef] [PubMed]
30. Koneru Neha. A Study on Covid associated Mucormycosis (CAM) with reference to Diabetic patients in Vizianagaram District, Andhra Pradesh, India, 2022. *International journal of advances in Engineering and Management.*, 4, 613-616. [Google Scholar] [CrossRef]
31. Cleveland Clinic. Pneumonia: Causes, Symptoms, Diagnosis & Treatment. *Health*, 2023 <https://my.clevelandclinic.org> [Google Scholar] [CrossRef]
32. Ashraf S. Ibrahim, Brad Spellberg, Thomas J. Walsh, Dimitrios P. Kontoyiannis, 2012 Pathogenesis of Mucormycosis, *Clinical Infectious Diseases.* 75. 16-22.
33. Mani Gudivada, M.Aruna, 2019- Dementia, a challenging health problem of elderly women in India- A Review, *International Journal of Research*, Volume VIII, Issue-V, PP 131-148.
34. Dr. Sankar Babu Gorla, Dr. Madhavi P, 2019- A cross sectional study on prescribing pattern on patients suffering from Chronic obstructive lung disease in a teaching hospital in Vizianagaram. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* Volume 18, Issue 1), PP 55-59.
35. Penny M Kris, William, S.Harris, Lawrence J. Apple. Fish consumption, Fish oil,  $\omega$ -3fatty acid and Cardio vascular disease, *Circulation, AHA journals*, [Google Scholar].
36. Kris-Etherton PM, Harris WS, Appel L et al., 2003 omega-3 fatty acids and cardiovascular

- disease, *Arteriosclerosis thrombosis and vascular biology*, 23(2)
37. Junichiro Tsuchiya et al., 2022 Epidemiological investigation of the factors affecting the COVID-19 case fatality rate, *Hirosaki Med.J.* 72:34-42
  38. Michael Alperovich, Mark I Neuman, Walter C Willett and Gary C Curhan, Fatty acid intake and Risk of Community acquired pneumonia in US Women., 2007 PMC online [Google Scholar]
  39. Cecilia A Hinojosa, Norberto Gonzalez Juarbe Md M Rahman, Gabriel Fernandes, Carlos J Orihuela and Marcos I Restrepo,  $\omega$ -3 fatty acid in contrast to  $\omega$ -6 fatty acid protection against pneumococcal pneumonia, PMC online [Google Scholar].
  40. Brigham, E. et al. The Johns Hopkins Post-Acute COVID-19 Team (PACT) 2021: a multidisciplinary, collaborative, ambulatory framework supporting COVID-19 survivors. *Am. J. Med.* <https://doi.org/10.1016/j.amjmed>.
  41. Hosey, M. M. & Needham, D. M. 2020. Survivorship after COVID-19 ICU stay. *Nat. Rev. Dis. Prim.* 6, 60.
  42. Carf, A., Bernabei, R., Landi, F. & Gemelli. 2020. Against COVID-19 Post-Acute Care Study Group. Persistent symptoms in patients after acute COVID-19. *J. Am. Med. Assoc.* 324, 603-605.
  43. Chopra, V., Flanders, S. A. 2020 Sixty-day outcomes among patients hospitalized with COVID-19. *Ann. Intern. Med.* <https://doi.org/10.7326/M20-5661>.
  44. Huang, C. et al. 2021 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet* 397, 220-232.