

REVIEW OF MEDICINE SURGERY AND DENTISTRY

VOLUME: 3 ISSUE: 1 YEAR: 2024

ISSN: 2957-417X



Cell Culture, Technology: Enhancing the Culture of Diagnosing Human Diseases

Shinde Pratiksha. A, Bangar Vilas B.

¹Dr. Vithalrao Vikhe Patil College of Pharmacy Viladghat Ahmednagar, India

²Ratnadeep College of Pharmacy, Ratnapur, Ahmednagar. Maharashtra, India

Abstract

The complicated processes of removing cells from their native environment (in vivo) and growing them in an artificially controlled environment (in vitro) are all part of cell culture. Because the availability of appropriate cell cultures is a prerequisite for the isolation of harmful viruses, cells from particular tissues or organs are cultivated as short-term or established cell lines that are frequently utilized for research and diagnostics, particularly in the context of viral infection. The necessary environment is provided by cell culture for the detection and identification of many human diseases. The "gold standard" for virus discovery is virus isolation in the cell culture. Researchers' opinions about the current application of cell culture technologies in the diagnosis.

Keywords: Pathogen discovery, Recombinant protein, Transgenic cell line, Viral isolation

INTRODUCTION

The early 20th century saw the development of the cell culture technique, which was used to examine animal cell behavior in vitro [1]. When embryologist Roux maintained chicken embryos in warm saline for several days, he developed the tissue culture concept, which laid the foundation for cell culture [2]. Thus, the term "cell culture" refers to the process of isolating animal cells and growing them in vitro in a controlled setting that is conducive to their growth [3, 4]. In order to achieve confluence, or the creation of a cell monolayer in a culture flask provided with the necessary nutrients and growth factors, this normally starts with a primary culture. After confluence is reached, the cells are either passed through or subculture from the necessary development factors

and nutrition. The cells are passaged or subculture from the primary to secondary and subsequence to tertiary until confluence is reached and a continuous cell line is created. It takes time and a lot of labor to isolate a virus in a cell culture [5, 6]. Many clinically significant viruses are still either difficult to grow in tissue culture or do not grow at all, while others may require a sophisticated culture technique that is either unavailable or not suited for use in diagnostic laboratories. These could lessen the usefulness of tissue culture in clinical diagnosis, which would make it less appealing for the diagnosis of human illnesses [5,7]. However, other scientists discovered that tissue culture is a reasonably objective method whose sole drawback is the virus's capacity to proliferate on the

chosen cell lines [8,9]. The ability to analyze a microorganism's genome without isolating the virus through cell culture has been made possible by recent developments in metagenomics and deep sequencing techniques. In order to finally identify the detected agent, this is accomplished using high-throughput sequencing employing a random amplified DNA product and comparing the sequences with a large bank of sequences. Because random primers can precisely amplify the template for sequencing without requiring prior knowledge about the suspected agent, this is made achievable [10–11]. In the field of pathogen detection, this approach is progressing quickly. It has been used indefinitely to find viruses, including the Schmallenberg, Bas-Congo, and Livia viruses [12, 13]. It is crucial to determine the infection's causal agent when dealing with critically ill individuals or outbreaks of infectious diseases. To identify the causal agent and recognize developing diseases, this study aims to describe some of the events in which viruses are isolated using molecular, serological, and electron microscope (EM) techniques, among other laboratory diagnosis assays.

METHODOLOGY

Utilizing the University Putra Malaysia online subscribed databases, searches for peer-reviewed journal articles in the field of health sciences and medicine were carried out utilizing search engines including Google Scholar, SCOPUS, and Medline. Except in cases where an older publication might be taken into consideration, all searches were restricted to those published between 2000 and 2015. Duplicates were eliminated, and all publications were in English. Articles published up until May 31, 2015, were the last ones that were searched. 2473 publications from the online database search were found; these were filtered based on the relevancy of the title and abstract, excluding conference abstracts, remarks, and brief messages, leaving 260 for full text review studies

Cell Culture and Electron Microscopy in Diagnosis

When identifying the cause of an uncommon clinical presentation, electron microscopy (EM) and cell culture separation play key roles. In a patient with a history of tick bites, the Bunya virus was isolated, according to one study [15]. Leukocytes from the patient who was initially suspected of having Ehrlichia pneumonia were inoculated into the DH82 cell line, which is a canine monocyte cell line, and the results showed some cytological alterations [Table/Fig-1]. The predicted bacterium was not found when the cells were processed for EM inspection; instead, a Bunya virus was detected. Bunya virus particles are detected as a bud in vesicles and extracellular areas of infected cells [Table/Fig-2]. With a little projection on the surface of the viral particles, the spherical virus envelope and the virus has granular core.

Cell Culture and RT-PCR

For the identification of influenza viruses, real-time reverse transcription polymerase chain reaction (qtr-PCR) and cell culture have been widely employed in clinical settings [16, 17]. However, it is labor-intensive, time-consuming, and requires highly skilled workers with specialized laboratory equipment and conditions, which makes it unsuitable for primary healthcare settings and low-income nations. Cell culture is still crucial for identifying the infection's causal agent in an outbreak, though. Cell culture and RTPCR were used to confirm the H7 N9 influenza cases that are now being reported [18].

Cell Culture

metabolomics Metabolomics in cell cultures is a useful tool for determining both the metabolic pathways that lead to the production of disease-related biomarkers and the biomarkers themselves. Through the discovery of new cancer biomarkers, metabolites are crucial in the diagnosis, recurrence, and prognosis of cancer. The construction of predictive models that will aid in the early detection of cancer is possible due to the

detection of a modest alteration in metabolism in the products of cellular processes.

Transgenic Cell Lines and Viral Detection

In cell culture, transgenic technology entails introducing stable genetic elements into the cell so that, upon infection, the virus induces the creation of readily detectable enzymes specific to the virus [19, 20]. But monoclonal antibody staining allows for its detection, and it can be found 16–24 hours after inoculation [21, 22]. On the other hand, a quicker transgenic system that can quickly identify HSV within 24 hours was created without the need for pricey monoclonal antibodies or medical training. The big rib nucleotide re-educates subunit is coded for by the HSV promoter derived from UL39 [23, 24].

Expression of Recombinant Protein for Detection of Influenza Virus Antibody

Recombinant protein technology has proven helpful for serological surveys of infections and is crucial in satisfying the need for quick, simple, and accurate tests in diagnostic laboratories [25]. It is possible to generate recombinant protein and utilize it to identify influenza virus antibodies.

SDS-PAGE was used to analyze the expressed protein's expression [Table/Fig-4]. Western blotting was then used to confirm the expression of the predicted 13 KDa protein, which was detected by polyclonal anti-NS antibody [26]. This demonstrated that Elisa, which offers a significant advantage over other methods for detecting particular antibodies, could be utilized to identify specific antibodies against influenza viruses utilizing the antigen. This demonstrated that Elisa could be used to identify certain antibodies against influenza viruses based on the antigen.

Issues arising from this review

uniformity in contrast to molecular techniques, the outcomes of cell culture might vary significantly based

on the methods used for collection, transportation, and handling [Table/Fig-3]. [Table/Fig. 4]:

the sample to preserve healthy inoculated cells and virus vitality [28]. Scholars engaged in debates on the value of cell culture in medical labs. While some people thought that there would be circumstances where using tissue culture in a diagnostic virology laboratory would be appropriate, others felt that this may not always be the case at the point of care, which would alter the role that cell culture plays in diagnostics [29, 30]. When necessary for a specific reason, viral isolation can be carried out by national laboratories with the necessary knowledge and cell culture systems, as well as by certain local regions [31].

Time-Consuming

In this day and age, when quick and precise clinical diagnosis is essential for prompt and efficient action, cell culture is quickly losing ground and importance in the diagnosis of human diseases. However, molecular methods offer a reliable and ageless means of diagnosis. As a result, molecular techniques are quickly replacing the conventional cell culture-based, early, and accurate diagnostic techniques, which have a substantial impact on patient care by limiting the extent of diseases through prompt treatment and, consequently, lowering the need for unnecessary hospital stays, the use of antibiotics, and the costs associated with them [32].

CONCLUSION


To sum up, cell culture is a vital technique in contemporary medicine with countless uses in the identification of human infections. The only restriction on cell culture techniques is the virus's capacity to proliferate in a certain cell line. But with the development of transgenic cell culture technologies, this has been resolved. Thus, we advise: Cell culture should be utilized to track the sensitivity and specificity of quick testing based on yearly antigen

assays, and the findings should be reported to the clinicians; Additionally, cell culture should be promoted for both positive results during low prevalence and negative fast test results received from patient features infection during high prevalence or outbreak: For the diagnosis of an unidentified virus, cell culture can also be utilized in conjunction with immunohistochemistry, histopathology, and PCR serological testing. Additionally, they help create a quick test for recently identified pathogens.

REFERENCES


- [1] Thorpe TA. History of plant tissue culture. *Molecular Biotechnology*. 2007;37(2):169-80.
- [2] Fung S, Wong F, Hussain M, Lok A. Sustained response after a 2-year course of lamivudine treatment of hepatitis B e antigen-negative chronic hepatitis B. *Journal of Viral Hepatitis*. 2004;11(5):432-38.
- [3] Ganem D, Schneider RJ. Hepadnaviridae: the viruses and their replication. *Fields Virology*. 2001; 2:2923-69.
- [4] Willmer EN. *Cells and tissues in culture: methods, Biology and Physiology*; Elsevier; 2013.
- [5] Leland DS, Ginocchio CC. Role of cell culture for virus detection in the age of technology. *Clinical Microbiology Reviews*. 2007;20(1):49-78.
- [6] Oberste MS, Nix WA, Maher K, Pallansch MA. Improved molecular identification of enteroviruses by RT-PCR and amplicon sequencing. *Journal of Clinical Virology*. 2003;26(3):375-77.
- [7] Griffith LG, Naughton G. Tissue engineering--current challenges and expanding opportunities. *Science*. 2002;295(5557):1009-14.
- [8] Khademhosseini A, Vacanti JP, Langer R. Progress in tissue engineering. *Scientific American*. 2009;300(5):64-71.
- [9] Ransohoff RM, Perry VH. Microglial physiology: unique stimuli, specialized responses. *Annual Review of Immunology*. 2009; 27:119-45.
- [10] Mokili JL, Rohwer F, Dutilh BE. Metagenomics and future perspectives virus discovery. *Current opinion in virology*. 2012;2(1):63-77
- [11] Lagier J-C, Edouard S, Pagnier I, Mediannikov O, Drancourt M, Raoult D. Current and past strategies for bacterial culture in clinical microbiology. *Clinical microbiology reviews*. 2015;28(1):208-36
- [12] Mehedi M. Ebola virus RNA editing: Characterization of the mechanism and gene products. 2011.
- [13] Liew FY, Pitman NI, McInnes IB. Disease-associated functions of IL-33: the new kid in the IL-1 family. *Nature Reviews Immunology*. 2010;10(2):103-10.
- [14] Deshmane SL, Kremlev S, Amini S, Sawaya BE. Monocyte chemoattractant protein-1 (MCP-1): an overview. *Journal of Interferon & Cytokine Research*. 2009;29(6):313-26.
- [15] Yu X-J, Liang M-F, Zhang S-Y, Liu Y, Li J-D, Sun Y-L, et al. Fever with thrombocytopenia associated with a novel bunyavirus in China. *New England Journal of Medicine*. 2011;364(16):1523-32
- [16] Templeton KE, Scheltinga SA, Beersma MF, Kroes AC, Claas EC. Rapid and sensitive method using multiplex real-time PCR for diagnosis of infections by influenza A and influenza B viruses, respiratory syncytial virus, and parainfluenza viruses 1, 2, 3, and 4. *Journal of Clinical Microbiology*. 2004;42(4):1564-69.

- [17] Rahman M, Vandermause MF, Kieke BA, Belongia EA. Performance of Binax NOW Flu A and B and direct fluorescent assay in comparison with a composite of viral culture or reverse transcription polymerase chain reaction for detection of influenza infection during the 2006 to 2007 season. *Diagnostic Microbiology and Infectious Disease*. 2008;62(2):162-66.
- [18] Xia J, Liu L, Wang L, Zhang Y, Zeng H, Liu P, et al. Experimental infection of pregnant rabbits with hepatitis E virus demonstrating high mortality and vertical transmission. *Journal of Viral Hepatitis*. 2015.
- [19] Deisseroth K, Airan RD. Cell line, system and method for optical control of secondary messengers. Google Patents; 2014.
- [20] Dimitrov DS. Virus entry: molecular mechanisms and biomedical applications. *Nature Reviews Microbiology*. 2004;2(2):109-22.
- [21] Anderson NW, Buchan BW, Ledebner NA. Light microscopy, culture, molecular, and serologic methods for detection of herpes simplex virus. *Journal of Clinical Microbiology*. 2014;52(1):2-8.
- [22] Atmar RL. Immunological Detection and Characterization. *Viral Infections of Humans*: Springer; 2014. pp. 47-62
- [23] Ma JZ, Russell TA, Spelman T, Carbone FR, Tschärke DC. Lytic gene expression is frequent in HSV-1 latent infection and correlates with the engagement of a cellintrinsic transcriptional response. *PLoS pathogens*. 2014;10(7): e1004237.
- [24] Fan F, Day S, Lu X, Tang Y-W. Laboratory diagnosis of HSV and varicella zoster virus infections. *Future Virology*. 2014;9(8):721-31.
- [25] Marks LV. *The Lock and Key of Medicine: Monoclonal Antibodies and the Transformation of Healthcare*: Yale University Press; 2015.
- [26] Schägger H. Tricine-SDS-PAGE. *Nature Protocols*. 2006; 1:16-22.
- [27] Portela A, Melero JA, Martínez C, Domingo E, Ortín J. A primer vector system that allows temperature dependent gene amplification and expression in mammalian cells: regulation of the influenza virus NSI gene expression. *Nucleic acids research*. 1985;13(22):7959-77.
- [28] Kao J-H, Chen P-J, Lai M-Y, Chen D-S. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology*. 2000;118(3):554-59.
- [29] Sacks G. Reproductive immunology: the relevance of laboratory research to clinical practice (and vice versa). *Human Reproduction*. 2014: deu325.
- [30] Viswanathan S, Keating A, Deans R, Hematti P, Prockop D, Stroncek DF, et al. Soliciting strategies for developing cell-based reference materials to advance mesenchymal stromal cell research and clinical translation. *Stem cells and development*. 2014;23(11):1157-67.
- [31] Stacey GN. *The Challenge of Standardization in Stem Cell Research and Development*. *Stem Cell Banking*: Springer; 2014. pp. 11-8.
- [32] Franco E, Bagnato B, Marino MG, Meleleo C, Serino L, Zaratti L. Hepatitis B: Epidemiology and prevention in developing countries. *World Journal of Hepatology*. 2012;4(3):74



[Figure-1]:

a) Visible virus-induced cytopathic effect in DH82 cells 8 days post-infection with Bunyavirus. The infected cells show visible granular particles and differentiate into macrophages with elongated pseudopodia. b) Bunya virus grown in Vero cells, detected on immunofluorescence assay [15].




[Figure-2]:

a) Negative stain Bunya virus purified from infected Vero cells. b) Transmission electron microscopy of virus-infected cells (DH82) shown by black arrows [15].



[Figure-3]

Result of successful positive recombinant colonies ligation. Lane 1: DNA ladder 1kb; Lane 2-13: positive colonies with NS1; Lane 4, 7, 8, and 12: negative colonies with no NS1 inserted [27]



[Figure-4]:

SDS-PAGE profile of un-purified expressed NS1 (13KDa) protein and un-induced plasmid. Lane 1: protein ladder; Lane 2 - 4: un-induced recombinant plasmid at zero hour; Lane 5 and 6: un-purified recombinant NS1 (13KDa) protein [26].